

BACKAID MAX Caplets

UPC #'s: 0-72959-53028-0, 0-72959-53038-9

Can I use BACKAID MAX Caplets with my prescription medication(s)?

You should ask your doctor or pharmacist if you can take **BACKAID MAX Caplets** while you are using other medications.

What is the dosage for BACKAID MAX Caplets?

The dosage for **BACKAID MAX** is two caplets with a full glass of water every 6 hours not to exceed 4 doses in 24 hours. Be sure to read all product package label Directions, Uses, and Warnings before use and follow them carefully.

What are the active ingredients in BACKAID MAX Caplets?

Each **BACKAID MAX Caplet** contains 500 mg acetaminophen, a pain reliever, and 25 mg of pamabrom, a mild diuretic.

Are BACKAID MAX Caplets gluten-free?

BACKAID MAX Caplets would qualify for the description of gluten-free as proposed by the FDA in that they do not contain any ingredients from wheat, rye, or barley and they contain less than 20 ppm gluten, as tested by a qualified independent laboratory.

Why are BACKAID MAX Caplets so difficult to remove from the blister package?

The difficulty you experienced opening **BACKAID MAX** blister packs is most likely attributable to the child resistant nature of the packs, commonly referred to as Child Resistant Closure (CRC) packaging. Certain packaging safety features are required by federal regulations on products such as **BACKAID MAX Caplets** and they may tend to reduce the ease of opening the package. We apologize for the inconvenience. You may find it is easier to open the blisters with scissors, as recommended on the blister back.