

URICALM Tablets

UPC #'s: 0-72959-41024-7, 0-72959-41036-0

What are the active ingredients in URICALM?

Each **URICALM** tablet contains 99.5mg of Phenazopyridine Hydrochloride.

What is the dosage for URICALM?

The dosage for **URICALM** is two tablets with water after meals as needed up to 3 times daily for 2 days maximum. Do not use more than 12 tablets in 2 days. Be sure to read all product package label Directions, Uses and Warnings and follow them carefully.

Are URICALM Tablets gluten-free?

URICALM Tablets would qualify for the description of gluten-free as proposed by the FDA in that they do not contain any ingredients from wheat, rye or barley and they contain less than 20 ppm gluten, as tested by a qualified independent laboratory.

Why can you use URICALM for only 2 days?

If you need to use **URICALM** for more than 2 days, that is an indication that the cause is not going to resolve itself and you need to see a doctor or other health care professional for further treatment.

It says that URICALM can interfere with laboratory tests. What tests are affected?

URICALM will discolor urine to a reddish-orange color, which may be of concern to the people administering the test. We do not have specific information to say which tests will be affected.

The label says this product may stain soft contact lenses. Does this mean do not handle soft contact lenses or do not wear soft contact lenses while you are using URICALM?

It means both. If you are concerned about staining, you should not wear soft contact lenses while using **URICALM**, and if you were to get the active ingredient on your fingers, you should not handle soft contact lenses until you have washed your hands thoroughly.