

Arthriten IPF Drug Facts (37/28-4RG, 0215)

Drug Facts	
Active ingredients (in each caplet)	Purpose
Acetaminophen 250 mg.....	Analgesic (pain reliever)
Aspirin 250 mg (NSAID*).....	Analgesic (pain reliever)
Caffeine Anhydrous 65 mg.....	Pain Reliever Aid
*Non-steroidal anti-inflammatory drug	
Uses For the temporary relief of minor aches and pains including the minor pain of arthritis.	
Warnings	
<p>Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock. ■ Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount for this product. ■ with other products containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.</p> <p>Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non-prescription NSAIDs [aspirin, ibuprofen, naproxen or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.</p>	
<p>Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor.</p>	
<p>Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, liver cirrhosis or kidney disease ■ you are taking a diuretic.</p>	
<p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout or arthritis ■ taking any other drug.</p>	
<p>Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ symptoms do not improve ■ pain or fever persists or gets worse ■ redness or swelling is present ■ ringing in the ears or hearing loss occurs ■ new symptoms occur ■ you experience any of the following signs of stomach bleeding: ■ you feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better.</p>	
<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	
<p>Directions ■ Read all package directions and warnings before use. Use only as directed. ■ Adults: Two (2) caplets every 6 hours while symptoms persist, not to exceed 4 doses (8 caplets) in 24 hours, or as directed by a doctor. ■ Drink a full glass of water with each dose ■ Intended for use by normally healthy adults only. ■ Persons under 18 years of age should use only as directed by a doctor.</p>	
<p>Other information **CONTENTS SEALED: Each ARTHRITEN white colored, capsule shaped caplet bears the identifying mark "ALVA" and a "2" on the reverse side, and is packaged in a bottle with a safety seal under the cap. Do not use if the seal appears broken or if product contents do not match product description. ■ You may report serious side effects to the phone number provided under <i>Questions?</i> below.</p>	
<p>Inactive ingredients Hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, silica, starch, stearic acid, talc, titanium dioxide and triglycerides.</p>	
<p>Questions? 1-800-792-2582</p>	